

References

- [1] Behrens, W.A. (1929). Ein Betrag zur Fehlenberechnung bei wenigen Beobachtungen. *Landwirtschaftliche Jahrbücher*, **68**, 807-837.
- [2] Box, G.E. and Tiao, G.C. (1973). Bayesian inference and statistical analysis. Reading, MA: Addison-Wesley.
- [3] Fisher, R.A. (1939). The comparison of samples with possibly unequal variance. *Ann. Eugenics*, **9**, 174-180.
- [4] Gelman, A., Carlin J., Stern, H. and Rubin, D. (1995). *Bayesian Data Analysis*. New York: Chapman and Hall.
- [5] Gelfand, A.E. and Smith, A.F.M. (1990). Sampling based approaches to calculating marginal densities. *J. Amer. Statist. Assoc.*, **85**, 398-409.
- [6] Lee, P.M. (1997). *Bayesian Statistics: An Introduction. 2nd edition*. London: Edward Arnold.
- [7] Lindley, D.V. and Smith, A.F.M. (1972). Bayes estimates for the linear model. *J. Roy. Statist. Soc. B*, **34**, 1-41.
- [8] Patil, V.H. (1965). Approximations to the Behrens-Fisher distribution. *Biometrika*, **52**, 267-271.
- [9] Spiegelhalter, D.J., Thomas, A. and Best, N.G. (1999). *WinBUGS Version 1.2 User Manual*. MRC Biostatistics Unit, Cambridge University.
- [10] Wiper, M.P. (1990). *Calibration and use of expert probability judgements*. Ph.D. thesis, School of Computer Studies, University of Leeds.
- [11] Wiper, M.P. and French, S. (1995). Combining experts' opinions using a normal-Wishart model. *Journal of Forecasting*, **14**, 25-34.
- [12] Wiper, M.P. and Pettit, L.I. (1996). On improving a model for combining experts' forecasts. In *Bayesian Statistics 5*, eds. J. M. Bernardo, J. O. Berger, A. P. Dawid y A. F. M. Smith. Oxford University Press, 809-813.