

References

- [1] Andreu, C., DeFreitas, N., Doucet, A. and Jordan, M. (2003). An introduction to MCMC for Machine Learning. *Machine Learning*, **50**, 5-43.
- [2] Bowmaker, J.K., Jacobs, G.H., Spiegelhalter, D.J. and Mollon, J.D. (1985). Two types of trichromatic squirrel monkey share a pigment in the red-green spectral region. *Vision Research*, **25**, 1937-1946.
- [3] Casella, G., Lavine, M. and Robert, C. (2000). Explaining the Perfect Sampler. *Research Report*, ISDS Duke University. Available from: <http://ftp.isds.duke.edu/WorkingPapers/00-16.html>
- [4] Chib, S. (1995). Marginal likelihood from the Gibbs output. *J. Amer. Statist. Assoc.*, **90**, 1313-1321.
- [5] Chib, S. and Greenberg, E. (1995). Understanding the Metropolis Hastings algorithm. *The American Statistician*, **49**, 327-335.
- [6] Gelfand, A.E. and Smith, A.F.M. (1990). Sampling based approaches to calculating marginal densities. *J. Amer. Statist. Assoc.*, **85**, 398-409.
- [7] Gelman, A., Carlin J., Stern, H. and Rubin, D. (1995). *Bayesian Data Analysis*. New York: Chapman and Hall.
- [8] Gelman, A. and Rubin, D.B. (1982). Inference from iterative simulation using multiple sequences. *Statistical Science*, **7**, 457-511.
- [9] Geman, S. and Geman, D. (1984). Stochastic relaxation, Gibbs distributions and the Bayesian restoration of images. *IEEE. Trans. Pattern Analysis and Machine Intelligence*, **6**, 721-741.
- [10] Gilks, W.R., Best, N.G. and Tan, K.K.C. (1995). Adaptive rejection Metropolis sampling within Gibbs sampling. *Applied Statist.*, **44**, 455-472.
- [11] Gilks, W.R., Richardson, S. and Spiegelhalter, D.J. (1996). *Markov chain Monte Carlo in Practice*. London: Chapman and Hall.
- [12] Grenander, U. and Miller, M. (1994). Representations of knowledge in complex systems. *J. Royal Statist. Soc. B*, **56**, 549-603.
- [13] Green, P.J. (1995). Reversible jump MCMC computation and Bayesian model determination. *Biometrika*, **82**, 711-732.
- [14] Hastings, W.K. (1970). Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, **57**, 97-109.
- [15] Mengerson, K. L., Robert, C. P. and Guihenneuc-Jouyaux, C. (1999). MCMC Convergence Diagnostics: A Review. In *Bayesian Statistics 6*, eds. J.M. Bernardo et al. Oxford: Oxford University Press.

- [16] Metropolis, N., Rosenbluth, A.W., Rosenbluth, M.N. and Teller, A.H. (1953). Equation of state calculations by fast computing machines. *J. Chem. Phys.*, **21**, 1087-1092.
- [17] Neal, R. (1996). *Bayesian learning for neural networks*. New York: Springer Verlag.
- [18] Neal, R. (2000). The slice sampler. *Research Report*, University of Toronto. Available from
<ftp://ftp.cs.utoronto.ca/pub/radford/mixmc.pdf>
- [19] Richardson, S. and Green, P. (1997). On Bayesian analysis of mixtures with an unknown number of components. *J. Royal Statist. Soc. B*, **59**, 731-792.
- [20] Robert, C.P. (1996). Inference in mixture models. In *Markov Chain Monte Carlo in Practice*, eds. Gilks et al. London: Chapman and Hall.
- [21] Robert, C.P. and Casella, G. (1999). *Monte Carlo Statistical Methods*. New York: Springer.
- [22] Roberts, G.O. and Smith, A.F.M. (1994). Simple conditions for the convergence of the Gibbs sampler and Metropolis Hastings algorithms. *Stochastic Processes and their Applications*, **49**, 207-216.
- [23] Roberts, G.O. and Tweedie, R.L. (1996). Geometric Convergence and Central Limit Theorems for Multidimensional Hastings and Metropolis Algorithms. *Biometrika*, **83**, 95-110.
- [24] Stephens, M. (2000). Bayesian analysis of mixtures with an unknown number of components — an alternative to reversible jump methods. *Ann. Statist.*, **28**, 40-74.
- [25] Tanner, M. and Wong, W. (1987). The calculation of posterior distributions by data augmentation. *J. Amer. Statist. Assoc.*, **82**, 528-550.